

AD-A167 345

EXPONENTIAL BOUNDS OF MEAN ERROR FOR THE KERNEL
ESTIMATES OF REGRESSION F. (U) PITTSBURGH UNIV PA
CENTER FOR MULTIVARIATE ANALYSIS L C ZHAO DEC 85


1/1

UNCLASSIFIED

TR-85-50 AFOSR-TR-86-0255 F49628-85-C-0008 F/G 12/1

NL






MICROCOPY

CHART

2


AFOSR-TR- 86 - 0 2 5 5

AD-A167 345



Center for Multivariate Analysis
University of Pittsburgh

DTIC FILE COPY



Approved for release by
the Defense Intelligence Agency

86 4 28 033

UNCLASSIFIED

AD-A162345

SECURITY CLASSIFICATION OF THIS PAGE

REPORT DOCUMENTATION PAGE

1a. REPORT SECURITY CLASSIFICATION UNCLASSIFIED			1b. RESTRICTIVE MARKINGS		
2a. SECURITY CLASSIFICATION AUTHORITY			3. DISTRIBUTION/AVAILABILITY OF REPORT Approved for public release; distribution unlimited.		
2b. DECLASSIFICATION/DOWNGRADING SCHEDULE			5. MONITORING ORGANIZATION REPORT NUMBER(S) AFOSR-TR. 86-0255		
4. PERFORMING ORGANIZATION REPORT NUMBER(S)			7a. NAME OF MONITORING ORGANIZATION Air Force Office of Scientific Research		
6a. NAME OF PERFORMING ORGANIZATION University of Pittsburgh		6b. OFFICE SYMBOL (If applicable)		7b. ADDRESS (City, State and ZIP Code) Directorate of Mathematical & Information Sciences, Bolling AFB DC 20332-6448	
6c. ADDRESS (City, State and ZIP Code) Dept Mathematics and Statistics Pittsburgh, PA 15260		8a. NAME OF FUNDING/SPONSORING ORGANIZATION AFOSR		8b. OFFICE SYMBOL (If applicable) NM	
8c. ADDRESS (City, State and ZIP Code) Bolling AFB DC 20332-6448		9. PROCUREMENT INSTRUMENT IDENTIFICATION NUMBER F49620-85-C-0008			
		10. SOURCE OF FUNDING NOS.			
		PROGRAM ELEMENT NO. 61102F		PROJECT NO. 2304	
				TASK NO. A5	
				WORK UNIT NO.	
11. TITLE (Include Security Classification) EXPONENTIAL BOUNDS OF MEAN ERROR FOR THE KERNAL ESTIMATES OF REGRESSION FUNCTIONS					
12. PERSONAL AUTHOR(S) L.C. ZHAO					
13a. TYPE OF REPORT Technical		13b. TIME COVERED FROM _____ TO _____		14. DATE OF REPORT (Yr., Mo., Day) 85 December	
15. PAGE COUNT 15					
16. SUPPLEMENTARY NOTATION					
17. COSATI CODES			18. SUBJECT TERMS (Continue on reverse if necessary and identify by block number)		
FIELD	GROUP	SUB. GR.			
19. ABSTRACT (Continue on reverse if necessary and identify by block number)					
<p>Let $(X, Y), (X_1, Y_1), \dots, (X_n, Y_n)$ be i.i.d. $R^T \times R$-valued random vectors with $E Y < \infty$, and let $Q_n(x)$ be a kernel estimate of the regression function $Q(x) = E(Y X = x)$. In this paper, we establish an exponential bound of the mean deviation between $Q_n(x)$ and $Q(x)$ given the training sample $Z^n = (X_1, Y_1, \dots, X_n, Y_n)$, under the conditions as weak as possible.</p>					
20. DISTRIBUTION/AVAILABILITY OF ABSTRACT UNCLASSIFIED/UNLIMITED <input checked="" type="checkbox"/> SAME AS RPT. <input type="checkbox"/> DTIC USERS <input type="checkbox"/>			21. ABSTRACT SECURITY CLASSIFICATION UNCLASSIFIED		
22a. NAME OF RESPONSIBLE INDIVIDUAL Major Brian W. Woodruff			22b. TELEPHONE NUMBER (Include Area Code) (202) 767-5027		22c. OFFICE SYMBOL NM

AIR FORCE OFFICE OF SCIENTIFIC RESEARCH (AFSC)
NOTICE OF TRANSMITTAL TO DTIC

This technical report is approved for release under the provisions of AFSC-12.
Distribution is unlimited.

MATTHEW J. KERPER

Chief, Technical Information Division

EXPONENTIAL BOUNDS OF MEAN ERROR FOR THE
KERNAL ESTIMATES OF REGRESSION FUNCTIONS

L. C. ZHAO

Center for Multivariate Analysis
University of Pittsburgh

Technical Report No. 85-50 ✓

December 1985

Center for Multivariate Analysis
University of Pittsburgh
515 Thackeray Hall
Pittsburgh, PA 15260

DTIC
ELECTE
MAY 01 1986
S D D

*Research sponsored by the Air Force Office of Scientific Research (AFSC) under Contract F49620-85-C-0008. The United States Government is authorized to reproduce and distribute reprints for governmental purposes notwithstanding any copyright notation herein.

ABSTRACT

Let $(X, Y), (X_1, Y_1), \dots, (X_n, Y_n)$ be i.i.d. $R^T \times R$ -valued random vectors with $E|Y| < \infty$, and let $Q_n(x)$ be a kernel estimate of the regression function $Q(x) = E(Y|X = x)$. In this paper, we establish an exponential bound of the mean deviation between $Q_n(x)$ and $Q(x)$ given the training sample $Z^n = (X_1, Y_1, \dots, X_n, Y_n)$, under the conditions as weak as possible.

1. INTRODUCTION

Let $(X, Y), (X_1, Y_1), \dots, (X_n, Y_n)$ be i.i.d. $R^r \times R$ -valued random vectors with $E|Y| < \infty$. To estimate $Q(x) = E(Y|X=x)$, the regression function of Y with respect to X , Nadaraya (1964) and Watson (1964) proposed a class of kernel estimates of the form

$$Q_n(x) = \frac{\sum_{i=1}^n K\left(\frac{X_i - x}{h}\right) Y_i}{\sum_{j=1}^n K\left(\frac{X_j - x}{h}\right)}, \quad (1)$$

where K is a probability density function on R^r , and $h=h_n$ is a sequence of positive numbers. Write

$$W_{ni}(x) = K\left(\frac{X_i - x}{h}\right) / \sum_{j=1}^n K\left(\frac{X_j - x}{h}\right), \quad (2)$$

we define $W_{ni}(x) = 1/n$, $i=1, 2, \dots, n$, when $0/0$ appears. Many scholars studied convergence problems of these estimates from various points of view. For the universal consistency, one can refer to, for example, Devroye and Wagner (1980I), Spiegelman and Sacks (1980). For the pointwise moment-consistency, see Devroye (1981). For the pointwise a.s. consistency, see Devroye (1981), Greblicki, Krzyzak and Pawlak (1984), Zhao and Fang (1985). In this paper, we study another convergence of these estimates.

Let $Z^n = (X_1, Y_1, \dots, X_n, Y_n)$ be a training sample, $g_n = g_n(x, Z^n)$ be an estimate of $Q(x)$. In some problems, we are interested in the following mean deviation of g_n given the training sample Z^n :

$$\begin{aligned} D(g_n) &= E\{|g_n(X, Z^n) - Q(X)| | Z^n\} \\ &= \int_{R^r} |g_n(x, Z^n) - Q(x)| F(dx), \end{aligned} \quad (3)$$

where F denotes the distribution of X . Devroye and Wagner (1980II) proved

that

$$\lim_{n \rightarrow \infty} D(Q_n) = 0 \quad \text{a.s.}$$

for the kernel estimates $Q_n(x)$ of $Q(x)$, if the following conditions are satisfied:

- (i) Y is bounded,
 - (ii) F has a density f ,
 - (iii) K is bounded and
- $$\int_{R^r} \psi(x) dx < \infty, \quad (4)$$

where

$$\psi(x) = \sup_{||u|| > ||x||} K(u), \quad x \in R^r$$

and $||\cdot||$ is the L_2 norm or L_∞ norm on R^r ,

- (iv) $h_n \rightarrow 0$ and $\sum_n \exp(-\alpha n h_n^r) < \infty$ for any $\alpha > 0$.

In this paper, we establish an exponential bound for the above mentioned deviation of Q_n . Take $||\cdot||$ as L_2 or L_∞ norm, and denote by $I(A)$ or I_A the indicator of set A . We establish the following

Theorem. Let $Q_n(x)$ be a kernel estimate defined by (1). Suppose that the following conditions are satisfied:

- (i) Y is bounded.
 - (ii) F , the distribution of X , has a density of f .
 - (iii) There exist positive constant α and ρ_0 such that
- $$K(x) \geq \alpha I(||x|| \leq \rho_0), \quad x \in R^r. \quad (5)$$

- (iv) $h \rightarrow 0$ and $nh^r \rightarrow \infty$ as $n \rightarrow \infty$.

Then for any given $\epsilon > 0$, we have

$$P\{D(Q_n) \geq \epsilon\} \leq e^{-cn}. \quad (6)$$

where $C > 0$ is a constant independent of n .

Obviously, we need only to give the proof for L_∞ norm. We shall introduce some lemmas in section 2, and give a proof of the theorem in section 3.

Accession For	
NTIS CRA&I	<input checked="" type="checkbox"/>
DTIC TAB	<input type="checkbox"/>
Unannounced	<input type="checkbox"/>
Justification	
By	
Distribution /	
Availability Codes	
Dist	Avail and/or Special
A-1	



2. SOME LEMMAS

For simplicity we use the following convention: $\varepsilon, \varepsilon_1, \varepsilon_2, \dots, C, C_0, C_1, \dots, \alpha, \beta_1, \beta_2, \delta$, ect., are all constants independent of n . $\#(A)$ denotes the cardinal of set A . $S_{x,\rho} = \{u \in \mathbb{R}^r : \|u-x\| \leq \rho\}$. F^* and λ^* denote the outer measure generated by F and the Lebesgue measure λ (on \mathbb{R}^r), respectively. F_n denotes the empirical measure of $X^n = (X_1, \dots, X_n)$. We now give four lemmas which are needed in the sequel.

Lemma 1 (Besicovitch Covering Lemma). Let E be a bounded subset of \mathbb{R}^r , and let K be a family of cubes covering E which contain a cube D_x with center x for each $x \in E$. Then there exist points $\{x_k\}$ in E such that

$$(i) \quad E \subset \bigcup D_{x_k},$$

(ii) there exists a constant σ depending only on d such that

$$\sum_k I(D_{x_k}) \leq \sigma.$$

For the proof, refer to Wheeden and Zygmund (1977), pp. 185-187.

Lemma 2. Let $T > 0$ be a given constant. Suppose that F has a density f . Then for any given $\varepsilon > 0$, we can choose $\beta_1 > 0$ small enough and $\beta_2 > 0$ large enough such that the set

$$E^* = \{x \in S_{0,T} : \beta_1(2\rho)^r \leq F(S_{x,\rho}) \leq \beta_2(2\rho)^r \quad (7)$$

for any $\rho \in (0,1)\}$

satisfies $F^*(S_{0,T} - E^*) < \varepsilon$.

Note that for any Borel-measurable set $E \subset E^*$, we have

$$\beta_1 \leq f(x) \leq \beta_2, \quad \text{for almost all } x \in E \text{ (with respect to } \lambda).$$

Proof. Set

$$E_1 = \{x \in S_{0,T}: \sup_{0 < \rho < 1} \lambda(S_{x,\rho}) / F(S_{x,\rho}) > 1/\beta_1\},$$

$$E_2 = \{x \in S_{0,T}: \sup_{0 < \rho < 1} F(S_{x,\rho}) / \lambda(S_{x,\rho}) > \beta_2\}.$$

For any $x \in E_1$ there exists a cube $S_{x,\rho}$ with $\rho \in (0,1)$ such that $\lambda(S_{x,\rho}) > F(S_{x,\rho}) > F(S_{x,\rho})/\beta_1$. By Lemma 1 there exist $x_k \in E_1$ and $S_{x_k,\rho_k} \triangleq S_k$ such that $\lambda(S_k) > F(S_k)/\beta_1$, $E_1 \subset \bigcup_k S_k$ and $\sum_k I(S_k) \leq \sigma$. Thus

$$\begin{aligned} F^*(E_1) &\leq F(\bigcup_k S_k) \leq \sum_k F(S_k) < \beta_1 \sum_k \lambda(S_k) \\ &= \beta_1 \int_{\bigcup_k S_k} I(S_k) d\lambda \leq \beta_1 \sigma \lambda(\bigcup_k S_k) \leq \beta_1 \sigma \lambda(S_{0,2T}). \end{aligned}$$

Taking $\beta_1 > 0$ small enough, we have $F^*(E_1) < \epsilon/2$. In the same way, we have $\lambda^*(E_2) \leq F(R^d)\sigma/\beta_2 = \sigma/\beta_2$. Taking β_2 large enough, we can make $\lambda^*(E_2)$ small enough and, by the absolute continuity of F with respect to λ , $F^*(E_2) < \epsilon/2$. The lemma is proved.

Fix $\delta \in (0, 1/2\sigma)$ and assume that $h = h_n \in (0,1)$. Set

$$G_n^* = \{x \in S_{0,T}: F_n(S_{x,h}) < \delta F(S_{x,h})\}. \quad (8)$$

Lemma 3. Suppose that F has a density f , $h = h_n \in (0,1)$ and $nn^r \rightarrow \infty$. Then for any given $\epsilon > 0$ we have

$$P\{F^*(G_n^*) \geq \epsilon\} < e^{-C_1 n}.$$

Proof. By Lemma 1, there exist $x_k \in G_n^*$ and $S_k = S_{x_k,h}$ such that $G_n^* \subset \bigcup_k S_k \triangleq G$, $\sum_k I(S_k) \leq \sigma$. Partition R^r into sets with the form $\prod_{j=1}^r [(i_j-1)eh, i_j eh]$, where $i_1, \dots, i_r = 0, \pm 1, \pm 2, \dots$ and e is a fixed constant to be chosen later. Call the partition ϕ , and write

$$\begin{aligned} \phi' &= \{B \in \phi: B \subset S_{0,2T}\}, \\ \tilde{G} &= \bigcup_{B \in G, B \in \phi} B, \quad \tilde{S}_{0,1} = \prod_{j=1}^r [-1+e, 1-e]. \end{aligned}$$

$$C_{x_k} \stackrel{\Delta}{=} S_k - \bigcup_{B \in \Phi, B \subset S_k} B \subseteq x_k + h(S_{0,1} - \bar{S}_{0,1}) \\ \stackrel{\Delta}{=} C_{x_k}^*.$$

Then

$$\lambda(C_{x_k}^*) = h^r \lambda(S_{0,1} - \bar{S}_{0,1}) = (2h)^r [1 - (1-e)^r] \\ \leq \text{re} \lambda(S_k).$$

Since

$$G - \tilde{G} \subset \bigcup_k (S_k - \bigcup_{B \in \Phi, B \subset S_k} B) \subset \bigcup_k C_{x_k}^*,$$

we see that

$$\lambda(G - \tilde{G}) \leq \sum_k \lambda(C_{x_k}^*) \leq \text{re} \sum_k \lambda(S_k) \\ \leq \text{re} \int_{US_k} I(S_k) d\lambda \leq \text{re} \sigma \lambda(\bigcup_k S_k) \\ \leq \text{re} \sigma \lambda(S_{0,2T}).$$

Hence we can choose e small enough to render $\lambda(G - \tilde{G})$ small enough and $F(G - \tilde{G}) < \varepsilon/4$. By (8) and the fact that $\sigma\delta < 1/2$, we get

$$F_n(\tilde{G}) \leq F_n(G) \leq \sum_k F_n(S_k) < \delta \sum_k F(S_k) \\ = \delta \int_{US_k} I(S_k) dF \leq \delta \sigma F(\bigcup_k S_k) = \delta \sigma F(G) \\ < \frac{1}{2} F(G).$$

Therefore

$$F(\tilde{G}) - F_n(\tilde{G}) > F(G) - \varepsilon/4 - \frac{1}{2} F(G) = \frac{1}{2} F(G) - \varepsilon/4,$$

and

$$F^*(G_n^*) \geq \varepsilon \quad \text{implies} \quad F(\tilde{G}) - F_n(\tilde{G}) > \varepsilon/4.$$

For any $H \subset \Phi'$, we write $UH = \bigcup_{B \in H} B$. Then

$$\{F^*(G_n^*) \geq \varepsilon\} \subset \bigcup_{H \subset \Phi'} \{F(UH) - F_n(UH) > \varepsilon/4\}.$$

Assume that $\varepsilon \in (0,1)$. By Hoeffding's inequality,

$$\sup_A P\{F(A) - F_n(A) > \varepsilon/4\} \leq \sup_A \exp\{-n(\varepsilon/4)^2/[2F(A) + \varepsilon/4]\}$$

$$\leq \exp\{-n\varepsilon^2/[16(2 + \varepsilon/4)]\} \leq \exp(-n\varepsilon^2/36).$$

Noticing that $\#\{H: H \subset \Phi'\} \leq 2^{C_0 h^{-r}}$ and $h^{-r} = o(n)$, we get

$$P\{F^*(G_n^*) \geq \varepsilon\} \leq 2^{C_0 h^{-r}} \sup_A P\{F(A) - F_n(A) \geq \varepsilon/4\}$$

$$\leq \exp(-C_1 n),$$

and the lemma is proved.

Lemma 4. Suppose that $\int_{\mathbb{R}^r} |g(x)|^p F(dx) < \infty$ for some $p > 0$, then

$$\lim_{h \rightarrow 0} \int_{S_{x,h}} |g(u) - g(x)|^p F(du) / F(S_{x,h}) = 0$$

for almost all x (with respect to F).

Refer to Wheeden and Zygmund (1977), p. 191, example 20.

3. PROOF OF THE THEOREM

Assume that $|Y| \leq M$. Without loss of generality, we can assume that $\rho_0 = 1$ in (5)(iii). It is enough to prove that for each fixed $T > 0$,

$$P\left\{\int_{S_{0,T}} |Q_n(x) - Q(x)| F(dx) > \varepsilon\right\} < e^{-cn}. \quad (9)$$

By Lemma 2, there exist $\beta_1 = \beta_1(\varepsilon)$, $\beta_2 = \beta_2(\varepsilon)$ and a compact set $E \subset E^*$ such that $F(S_{0,T} - E) < \varepsilon/8M$, where E^* is defined by (7). Hence

$$\int_{S_{0,T} - E} |Q_n(x) - Q(x)| F(dx) \leq 2MF(S_{0,T} - E) < \varepsilon/4.$$

Fix $\delta \in (0, 1/2\sigma)$. By Lemma 3, there exists a compact set H_n such that

$$H_n \subset \{x \in S_{0,T} : F_n(S_{x,h}) \geq \delta F(S_{x,h})\}, \quad (10)$$

and

$$P\{F(S_{0,T} - H_n) \geq \varepsilon/(8M)\} < e^{-C_1 n} \quad (11)$$

Hence

$$P\left\{\int_{S_{0,T} - H_n} |Q_n(x) - Q(x)| F(dx) > \varepsilon/4\right\} < e^{-C_1 n}.$$

Therefore, we need only to prove that

$$P\left\{\int_{H_n \cap E} |Q_n(x) - Q(x)| F(dx) > \varepsilon/2\right\} < e^{-C_2 n}. \quad (12)$$

For $x \in H_n \cap E$, by (5)(iii), (7) and (10), we have

$$\begin{aligned} \bigwedge_{j=1}^n K\left(\frac{X_j - x}{h}\right) &\geq n\alpha F_n(S_{x,h}) \geq n\alpha\delta F(S_{x,h}) \\ &\geq n\alpha\delta\beta_1 2^{r_r} h^r, \end{aligned}$$

and $f(x) \leq \beta_2$. Write $C_3 = \beta_2/(\alpha\delta\beta_1 2^{r_r})$, we see that

$$\begin{aligned} \int_{H_n \cap E} |Q_n(x) - Q(x)| f(x) dx \\ \leq C_3 (nh^r)^{-1} \int_{H_n \cap E} \bigwedge_{i=1}^n K\left(\frac{X_i - x}{h}\right) (Y_i - Q(x)) dx. \end{aligned}$$

There exist finite positive constants m, a_1, \dots, a_m and disjoint regular cubes A_1, \dots, A_m such that $K^*(x) = \sum_{i=1}^m a_i I_{A_i}(x)$ satisfies

$$\int_{R^r} |K(x) - K^*(x)| dx < \varepsilon / (8C_3M).$$

Here a regular cube means a r -fold product of one-dimensional compact intervals.

Thus

$$\begin{aligned} P\{(nh^r)^{-1} \int_{H_n \cap E} \left| \sum_{i=1}^n \left(K\left(\frac{X_i - x}{h}\right) - K^*\left(\frac{X_i - x}{h}\right) \right) (Y_i - Q(x)) \right| dx \\ \leq 2C_3M(nh^r)^{-1} \sum_{i=1}^n \int \left| K\left(\frac{X_i - x}{h}\right) - K^*\left(\frac{X_i - x}{h}\right) \right| dx \\ \leq 2C_3M \int |K(x) - K^*(x)| dx < \varepsilon/4. \end{aligned}$$

Take $\varepsilon_1 = \varepsilon / (4C_3)$. To prove (12), it is enough to prove that

$$\begin{aligned} P\{(nh^r)^{-1} \int_{H_n \cap E} \left| \sum_{i=1}^n K\left(\frac{X_i - x}{h}\right) (Y_i - Q(x)) \right| dx > \varepsilon_1\} \\ < e^{-C_2n}. \end{aligned}$$

It is sufficient for any $\varepsilon_2 > 0$ and any regular cube A to prove that

$$\begin{aligned} P\{(nh^r)^{-1} \int_{H_n \cap E} \left| \sum_{i=1}^n I_{x+hA}(X_i) (Y_i - Q(X_i)) \right| dx \geq \varepsilon_2\} \\ < e^{-C_3n}, \end{aligned} \quad (13)$$

and

$$\begin{aligned} P\{(nh^r)^{-1} \int_{H_n \cap E} \left| \sum_{i=1}^n I_{x+hA}(X_i) (Q(X_i) - Q(x)) \right| dx \geq \varepsilon_2\} \\ < e^{-C_3n}. \end{aligned} \quad (14)$$

We proceed to prove (13). To this end, we construct the partition Φ of R^r mentioned in the proof of Lemma 3. Assume that $A = \prod_{i=1}^r [x_i, x_i + a_i]$ and $\min a_i \geq 2e$. Set $\tilde{A} = \prod_{i=1}^r [x_i + e, x_i + a_i - e]$. $A_x = \bigcup_{B \in \Phi, B \subseteq x+hA} B$.

$$C_x = x + hA - A_x x + h(A - \bar{A}) = C_x^*.$$

It is easy to see that we can make $\lambda(A - \bar{A})$ arbitrarily small by choosing h small enough. We have

$$\begin{aligned} & (nh^r)^{-1} \int_{H_n \cap E} \left| \sum_{i=1}^n I_{x+hA}(X_i)(Y_i - Q(X_i)) \right| dx \\ & \leq (nh^r)^{-1} \int_{H_n \cap E} \left| \sum_{i=1}^n I_{A_x}(X_i)(Y_i - Q(X_i)) \right| dx \\ & \quad + 2Mh^{-r} \int_{H_n} (C_x^*) dx \\ & \leq (nh^r)^{-1} \int_{H_n \cap E} \left| \sum_{i=1}^n I_B(X_i)(Y_i - Q(X_i)) \right| dx \\ & \quad + 2M\lambda(A - \bar{A}). \end{aligned} \tag{15}$$

Here we use the fact that $\int v(x+hD)dx = h^r \lambda(D)$ for any r -dimensional probability measure v and any Borel set $D \subset \mathbb{R}^r$. We can choose h such that $2M\lambda(A - \bar{A}) < \varepsilon_2/2$. Note that for $B \in \Phi$, $\lambda\{x, B \subseteq x+hA\} \leq C_4 h^r$, and $\bigcup_{x \in H_n \cap E} \{x+hA\} \subset S_{0,2T}$ for small h .

Hence, for large n , we have

$$(15) \leq C_4 n^{-1} \int_{B \in \Phi} \left| \sum_{i=1}^n I_B(X_i)(Y_i - Q(X_i)) \right| + \varepsilon_2/2.$$

Set $\varepsilon_3 = \varepsilon_2/(4C_4)$. To prove (13), we need only to prove that

$$P\left\{ \sum_{B \in \Phi} \left| \sum_{i=1}^n I_B(X_i)Y_i - n \int_B Q(x) dF \right| \geq n\varepsilon_3 \right\} < e^{-C_5 n}, \tag{16}$$

$$P\left\{ \sum_{B \in \Phi} \left| \sum_{i=1}^n I_B(X_i)Q(X_i) - n \int_B Q(x) dF \right| \geq n\varepsilon_3 \right\} < e^{-C_5 n}. \tag{17}$$

Let N be a Poisson random variable with mean value n , which is independent of $(X_1, Y_1), (X_2, Y_2), \dots$. In the sequel, we use \sum' for $\sum_{B \in \Phi}$. Notice

That B 's are disjoint, we see that for $t_B \in (-\infty, \infty)$,

$$\begin{aligned} & E\left\{\exp\left[\sum_{i=1}^N t_B I_B(X_i) Y_i\right]\right\} \\ &= \sum_{\ell=0}^{\infty} \frac{n^\ell e^{-n}}{\ell!} [E\{\exp\left[\sum_{i=1}^N t_B I_B(X_i) Y_i\right]\}]^\ell \\ &= \exp\left\{n \sum_{i=1}^N E[I_B(X_i) (e^{t_B Y_i} - 1)]\right\}, \end{aligned}$$

So that, $\{|\sum_{i=1}^N I_B(X_i) Y_i - n \int_B Q(x) dF|, B \in \Phi\}$ is a group of mutually independent variables. Set

$$Z(B, N) = \sum_{i=1}^N I_B(X_i) Y_i - n \int_B Q(x) dF.$$

for $t > 0$, notice that $e^t - t \geq e^{-t} + t$, we have

$$\begin{aligned} P\left\{\sum_{i=1}^N |Z(B, N)| \geq \frac{1}{2} n \epsilon_3\right\} &\leq \exp(-\frac{1}{2} t n \epsilon_3) E \exp\left\{t \sum_{i=1}^N |Z(B, N)|\right\} \\ &= \exp(-\frac{1}{2} t n \epsilon_3) \prod_{B \in \Phi} E\{\exp(t |Z(B, N)|)\} \\ &\leq \exp(-\frac{1}{2} t n \epsilon_3) \prod_{B \in \Phi} [E \exp(t Z(B, N)) + E \exp(-t Z(B, N))] \\ &= \exp(-\frac{1}{2} t n \epsilon_3) \prod_{B \in \Phi} \{ \exp[n E I_B(X_1) (e^{t Y_1} - 1)] \\ &\quad + \exp[n E I_B(X_1) (e^{-t Y_1} - 1)] \} \\ &\leq \exp(-\frac{1}{2} t n \epsilon_3) \prod_{B \in \Phi} \{ 2 \exp[n E I_B(X_1) (e^{t |Y_1|} - t |Y_1| - 1)] \} \\ &\leq \exp(-\frac{1}{2} t n \epsilon_3) 2^{C_0 h^{-r}} \exp\left\{n \sum_{i=1}^N E[I_B(X_i) (e^{t M} - t M - 1)]\right\} \\ &\leq \exp(-\frac{1}{2} t n \epsilon_3) e^{o(n)} \exp(n(e^{t M} - t M - 1)). \end{aligned} \tag{18}$$

Take $t \in (0, 1/M)$, we see that $e^{t M} - t M - 1 \leq t^2 M^2$. Hence, we can take $t > 0$ such that

$$P\left\{\sum_{i=1}^N \left| \sum_{i=1}^N I_B(X_i) Y_i - n \int_B Q(x) dF \right| \geq \frac{1}{2} n \epsilon_3\right\} < e^{-C_6 n}. \tag{19}$$

Write $\Delta = (X_1, X_2, \dots)$. By Jensen's inequality, for $t > 0$ we have

$$\begin{aligned}
 & P\left\{\left|\sum_{i=1}^N I_B(X_i)Q(X_i) - n \int_B Q(x)dF\right| \geq \frac{1}{2}n\epsilon_3\right\} \\
 &= P\left\{\left|E(Z(B,N)|\Delta)\right| \geq \frac{1}{2}n\epsilon_3\right\} \\
 &\leq P\left\{E\left(\sum_{i=1}^N |Z(B,N)| |\Delta\right) \geq \frac{1}{2}n\epsilon_3\right\} \\
 &\leq \exp(-\frac{1}{2}t n \epsilon_3) E\left\{\exp\left(t E\left(\sum_{i=1}^N |Z(B,N)| |\Delta\right)\right)\right\} \\
 &\leq \exp(-\frac{1}{2}t n \epsilon_3) E\left\{E\left[\exp\left(t \sum_{i=1}^N |Z(B,N)|\right) |\Delta\right]\right\} \\
 &= \exp(-\frac{1}{2}t n \epsilon_3) E\left[\exp\left(t \sum_{i=1}^N |Z(B,N)|\right)\right].
 \end{aligned}$$

By (18) and (19), we can take $t > 0$ such that

$$P\left\{\left|\sum_{i=1}^N I_B(X_i)Q(X_i) - n \int_B Q(x)dF\right| \geq \frac{1}{2}n\epsilon_3\right\} < e^{-C_6 n}. \quad (20)$$

Note that

$$\begin{aligned}
 \left|\sum_{i=1}^N I_B(X_i)Y_i - \sum_{i=1}^n I_B(X_i)Y_i\right| &\leq M|N-n|, \\
 \left|\sum_{i=1}^N I_B(X_i)Q(X_i) - \sum_{i=1}^n I_B(X_i)Q(X_i)\right| &\leq M|N-n|,
 \end{aligned}$$

we have

$$P\left\{\left|\sum_{i=1}^N I_B(X_i)Y_i - \sum_{i=1}^n I_B(X_i)Y_i\right| \geq \frac{1}{2}n\epsilon_3\right\} \quad (21)$$

$$\leq P\{|N-n| \geq n\epsilon_3/(2M)\} < e^{-C_7 n},$$

$$\begin{aligned}
 & P\left\{\left|\sum_{i=1}^N I_B(X_i)Q(X_i) - \sum_{i=1}^n I_B(X_i)Q(X_i)\right| \geq \frac{1}{2}n\epsilon_3\right\} \\
 & < e^{-C_7 n}.
 \end{aligned} \quad (22)$$

From (19) - (22), (16) and (17) follows, and (13) is proved. It remains to prove (14). To this end, we need only to prove

$$P\left\{\sum_{i=1}^n Z(X_i) \geq n\epsilon_2\right\} < e^{-C_3 n}, \quad (23)$$

where

$$\begin{aligned} Z(u) &= h^{-r} \int_{R^r} I_{x+hA}(u) |Q(u) - Q(x)| dx \\ &\leq 2Mh^{-r} \int_A \left(\frac{u-x}{h}\right) dx = 2M\lambda(A) \triangleq C_8 \end{aligned} \quad (24)$$

Hence, we have

$$\left| \sum_{i=1}^N Z(X_i) - \sum_{i=1}^n Z(X_i) \right| \leq C_8 |N-n|,$$

and

$$\begin{aligned} P\left\{\sum_{i=1}^N Z(X_i) - \sum_{i=1}^n Z(X_i) \geq \frac{1}{2}n\epsilon_2\right\} \\ \leq P\{|N-n| \geq n\epsilon_2/(2C_8)\} < e^{-C_9 n}. \end{aligned} \quad (25)$$

For $t > 0$, we have

$$\begin{aligned} P\left\{\sum_{i=1}^N Z(X_i) > \frac{1}{2}n\epsilon_2\right\} &\leq \exp(-\frac{1}{2}t n\epsilon_2) E\left\{\exp\left(t \sum_{i=1}^N Z(X_i)\right)\right\} \\ &= \exp\left(-\frac{1}{2}t n\epsilon_2 + n \int (e^{tZ(u)} - 1) F(du)\right). \end{aligned} \quad (26)$$

Take $t \in (0, 1/C_8)$. By $0 \leq tZ(u) \leq 1$ we get

$$n \int (e^{tZ(u)} - 1) F(du) \leq 2nt \int Z(u) F(du) \quad (27)$$

Take $\rho > 0$ so large that $A \subset S_{0,\rho}$. Then, by Lemma 4, we have

$$\begin{aligned} Z(u) &= h^{-r} \int_{u-hA} |m(x) - m(u)| dx \\ &\leq \lambda(S_{0,\rho}) \int_{S_{u,h\rho}} |m(x) - m(u)| dx / \lambda(S_{u,h\rho}) \end{aligned}$$

$\rightarrow 0$ as $h \rightarrow 0$, for almost all $x(\lambda)$.

In view of (24), from the dominated convergence theorem, we see that

$$\lim_{h \rightarrow 0} \int Z(u) F(du) = 0. \quad (28)$$

By (26) - (28), we can take $t > 0$ sufficiently small such that

$$\begin{aligned} P\left\{ \sum_{i=1}^N Z(X_i) > \frac{1}{2} n \varepsilon_2 \right\} &\leq \exp(-\frac{1}{2} n t \varepsilon_2 + o(nt)) \\ &< e^{-C_{10} n}. \end{aligned}$$

From (25) and (29), we obtain (23), and (14) follows. Up to now, the theorem is proved.

REFERENCES

- [1] Devroye, L.P. (1981). On the Almost Everywhere Convergence of Non-parametric Regression Function Estimates. Ann. Statist., 9, 1310-1319.
- [2] Devroye, L.P. and Wagner, T.J. (1980I). Distribution Free Consistency Results in Nonparametric Discrimination and Regression Function Estimation. Ann. Statist., 8, 231-239.
- [3] Devroye, L.P. and Wagner, T.J. (1980II). On the L_1 Convergence of Kernel Estimators of Regression Functions With Applications in Discrimination. Z. Wahrscheinlichkeitstheorie und Verw. Gebiete, 51, 15-25.
- [4] Greblicki, W., Krzyak, A. and Pawlak, M. (1984). Distribution-free Pointwise Consistency of Kernel Regression Estimate. Ann. Statist., 12, 1570-1575.
- [5] Nadaraya, E.A. (1964). On Estimating Regression. Theor. Probability and Appl., 9, 141-142.
- [6] Spiegelman, C. and Sacks, J. (1980). Consistent Window Estimation in Nonparametric Regression. Ann. Statist., 8, 240-246.
- [7] Watson, G.S. (1964). Smooth Regression Analysis. Sankhya, Ser. A, 26, 359-372.
- [8] Wheeden, R.L. and Zygmund, A. Measure and Integral, Marcel Dekker, New York, 1977.
- [9] Zhao, Lincheng and Fang, Zhaoben (1985). Strong Convergence of Kernel Estimates of Nonparametric Regression Functions. Chin. Ann. of Math., Ser. B, 6, 147-155.

END

FILMED

6-86

DTIC